



GoldBio Competent Cells Characteristics and Applications

Products	Transformation Efficiency	Applications and Characteristics
<u>BL21 (DE3) Chemically Competent E. coli Cells (CC-103)</u>	$\geq 5 \times 10^5$ CFU/ μ g	– Routine protein expression from non-T7 vectors – For routine T7 expression – Deficient in Lon and OmpT proteases/B strain – Resistant to phage T1 (<i>fhuA2</i>)
<u>BL21 (DE3) Electrocompetent E. coli Cells (CC-204)</u>	$\geq 1 \times 10^8$ CFU/ μ g	– Protein expression and complex library expression – For routine T7 expression – Deficient in Lon and OmpT proteases – Resistant to phage T1 (<i>fhuA2</i>) – B strain
<u>BL21 Chemically Competent E. coli Cells (CC-102)</u>	$\geq 1 \times 10^6$ CFU/ μ g	– Routine protein expression from non-T7 vectors – Deficient in Lon and OmpT proteases – Resistant to phage T1 (<i>fhuA2</i>)

GoldBio Competent Cells Characteristics and Applications



Products	Transformation Efficiency	Applications and Characteristics
<u>CJ236 Chemically Competent E. Coli Cells (CC-151)</u>	$\geq 1 \times 10^4$ CFU/ μ g	– Kunkel method of site-directed mutagenesis – dUTPase deficiency – Uracil-DNA glycosylase deficiency – Thiamine auxotrophy
<u>CJ236 Electrocompetent E. Coli Cells (CC-251)</u>	$\geq 1 \times 10^9$ CFU/ μ g	– Kunkel method of site-directed mutagenesis – dUTPase deficiency – Uracil-DNA glycosylase deficiency – Thiamine auxotrophy



GoldBio Competent Cells Characteristics and Applications

Products	Transformation Efficiency	Applications and Characteristics
<u>DL39 (DE3) Chemically Competent E. coli Cells (CC-104)</u>	$\geq 1 \times 10^7$ CFU/ μ g	– Transformation and protein expression – Deficient in aromatic, branched-chain and aspartate transaminases – For routine T7 expression
<u>GB10B-Pro™ Electrocompetent E. coli Cells (CC-201)</u>	$\geq 1 \times 10^9$ CFU/ μ g	– Cloning synthetic bio-applications, BAC cloning, assembling large and multi-DNA fragments

GoldBio Competent Cells Characteristics and Applications



Products	Transformation Efficiency	Applications and Characteristics
<u>GB10B™ Chemically Competent E. coli Cells (CC-100)</u>	$\geq 8.2 \times 10^6$ CFU/ μ g	– Cloning and subcloning – Blue/white screening ($\phi 80\text{lacZ}\Delta\text{M15}$) – <i>mcrA</i> <i>mcrBC</i>, and <i>mrr</i> deletion for cloning of methylcytosine and methyladenine-containing DNA – Endonuclease deficient (<i>endA1</i>)
<u>GB10B™ Electrocompetent E. coli Cells (CC-200)</u>	$\geq 2 \times 10^8$ CFU/ μ g	– Blue/white screening ($\phi 80\text{lacZ}\Delta\text{M15}$) – <i>mcrA</i> <i>mcrBC</i>, and <i>mrr</i> deletion for cloning of methylcytosine and methyladenine-containing DNA – High transformation efficiency
<u>GB5-alpha™ Chemically Competent E. coli Cells (CC-101)</u>	$\geq 1 \times 10^8$ CFU/ μ g	– Cloning and subcloning – Blue/white screening ($\phi 80\text{lacZ}\Delta\text{M15}$) – High-efficiency transformation of recombinant DNA – Cloning guide RNAs (gRNAs) for CRISPR applications – Amplification of plasmids after site-directed mutagenesis – Propagation of synthetic biology constructs and genetic circuits

GoldBio Competent Cells Characteristics and Applications

Products	Transformation Efficiency	Applications and Characteristics
<u>GB5-alpha™ Electrocompetent E. coli Cells (CC-203)</u>	$\geq 5 \times 10^8$ CFU/ μ g	– High efficiency transformation for many applications including cloning and subcloning – Increased plasmid yield and improved plasmid quality (<i>endA1</i> and <i>recA1</i> mutations)
<u>HB101 Chemically Competent E. coli Cells (CC-150)</u>	$\geq 7 \times 10^6$ CFU/ μ g	– Cloning and subcloning – non Blue/white screening – Prevents cleavage of cloned DNA by endogenous restriction enzymes (or the <i>hsdS20</i>(rB-mB-) restriction minus genotype) – Contain the <i>recA13</i> mutation that minimizes recombination and helps insert stability
<u>JM109 Chemically Competent E. coli Cells (CC-114)</u>	$\geq 6 \times 10^5$ CFU/ μ g	– Cloning and subcloning – Library construction – Plasmid isolation – Blue-white screening

GoldBio Competent Cells Characteristics and Applications

Products	Transformation Efficiency	Applications and Characteristics
<u>RR1 Chemically Competent E. coli Cells (CC-113)</u>	$\geq 3 \times 10^7$ CFU/ μ g	– Cloning and subcloning – recA+ derivative of the HB101 strain
<u>TG1 Phage Display Electrocompetent Cells (CC-205)</u>	$\geq 1 \times 10^9$ CFU/ μ g	– Protein expression – Amber suppressor strain (<i>supE</i>) – Phage display library screening
<u>AGL-1 Agrobacterium Chemically Competent Cells (CC-106)</u>	$\geq 4 \times 10^4$ CFU/ μ g	– High transformation efficiency – T-DNA binary system with the Ti plasmid pTiBO542 – Rifampicin and carbenicillin resistance

GoldBio Competent Cells Characteristics and Applications

Products	Transformation Efficiency	Applications and Characteristics
<u>AGL-1 Agrobacterium Electrocompetent Cells (CC-208)</u>	$\geq 5 \times 10^7$ CFU/ μ g	– High transformation efficiency – cDNA or gDNA library construction – T-DNA binary system with the Ti plasmid pTiBO542 – Rifampicin and carbenicillin resistance
<u>AGL-1 (pSoup) Agrobacterium Electrocompetent Cells (CC-218)</u>	$\geq 2 \times 10^8$ CFU/ μ g	– High transformation efficiency – cDNA or gDNA library construction – T-DNA binary system with the Ti plasmid pTiBO542 – Rifampicin and carbenicillin resistance
<u>AGL-1 (pSoup) Agrobacterium Chemically Competent Cells (CC-116)</u>	$\geq 3 \times 10^3$ CFU/ μ g	– High transformation efficiency – cDNA or gDNA library construction – Stabilizes recombinant plasmids (<i>recA</i> mutation) – T-DNA binary system with the Ti plasmid pTiBO542 – Rifampicin and carbenicillin resistance

GoldBio Competent Cells Characteristics and Applications



Products	Transformation Efficiency	Applications and Characteristics
<u>AGL-1 (pSoup-P19) Agrobacterium Electrocompetent Cells (CC-228)</u>	$\geq 1 \times 10^8$ CFU/ μ g	– High transformation efficiency – cDNA or gDNA library construction – Stabilizes recombinant plasmids (<i>recA</i> mutation) – T-DNA binary system with the Ti plasmid pTiBO542 – Rifampicin and carbenicillin resistance
<u>AGL-1 (pSoup-P19) Agrobacterium Chemically Competent Cells (CC-126)</u>	$\geq 2 \times 10^3$ CFU/ μ g	– High transformation efficiency – cDNA or gDNA library construction – Stabilizes recombinant plasmids (<i>recA</i> mutation) – T-DNA binary system with the Ti plasmid pTiBO542 – Rifampicin and carbenicillin resistance
<u>AGL-1 (pSuperAgro™ v4) Agrobacterium Electrocompetent Cells (CC-624)</u>	$\geq 1 \times 10^7$ CFU/ μ g	– High transformation efficiency – AcdS and GabT activity, driven by a single lac promotor – cDNA or gDNA library construction – Stabilizes recombinant plasmids (<i>recA</i> mutation) – T-DNA binary system with the Ti plasmid pTiBO542 – Rifampicin and carbenicillin resistance

GoldBio Competent Cells Characteristics and Applications



Products	Transformation Efficiency	Applications and Characteristics
<u>AGL-1 (pSuperAgro™ v4) Agrobacterium Chemically Competent Cells (CC-524)</u>	$\geq 1 \times 10^3$ CFU/ μ g	– High transformation efficiency – AcdS and GabT activity, driven by a single lac promotor – cDNA or gDNA library construction – Stabilizes recombinant plasmids (<i>recA</i> mutation) – T-DNA binary system with the Ti plasmid pTiBO542 – Rifampicin and carbenicillin resistance
<u>C58C1 Agrobacterium Chemically Competent Cells (CC-109)</u>	$\geq 1 \times 10^5$ CFU/ μ g	– Very high transformation efficiency – T-DNA binary system with the Ti plasmid pTiC58 – Rifampicin resistance and streptomycin resistance – <i>Agrobacterium</i>-mediated transformation
<u>C58C1 Agrobacterium Electrocompetent Cells (CC-240)</u>	$\geq 1 \times 10^7$ CFU/ μ g	– Very high transformation efficiency – T-DNA binary system with the Ti plasmid pTiC58 – Rifampicin resistance and streptomycin resistance – <i>Agrobacterium</i>-mediated transformation
<u>C58C1 (pSuperAgro™ v4) Agrobacterium Electrocompetent Cells (CC-654)</u>	$\geq 1 \times 10^7$ CFU/ μ g	– Very high transformation efficiency – cDNA or gDNA library construction – T-DNA binary system with the Ti plasmid pTiC58 – Rifampicin resistance and streptomycin resistance – <i>Agrobacterium</i>-mediated transformation

GoldBio Competent Cells Characteristics and Applications



Products	Transformation Efficiency	Applications and Characteristics
<u>C58C1 (pSuperAgro™ v4) Agrobacterium Chemically Competent Cells (CC-554)</u>	$\geq 1 \times 10^3$ CFU/ μ g	– Very high transformation efficiency – cDNA or gDNA library construction – T-DNA binary system with the Ti plasmid pTiC58 – Rifampicin resistance and streptomycin resistance – <i>Agrobacterium</i>-mediated transformation
<u>EHA105 Agrobacterium Chemically Competent Cells (CC-108)</u>	$\geq 5 \times 10^4$ CFU/ μ g	– High transformation efficiency – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pEHA105(pTiBo542DT-DNA) – Rifampicin resistance
<u>EHA105 Agrobacterium Electrocompetent Cells (CC-225)</u>	$\geq 1.6 \times 10^8$ CFU/ μ g	– High transformation efficiency – cDNA or gDNA library construction – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pEHA105(pTiBo542DT-DNA) – Rifampicin resistance (pTiBo542DT-DNA)
<u>Auxo-Agro® EHA105 Electrocompetent Cells (CC-268)</u>	$\geq 1.6 \times 10^8$ CFU/ μ g	– High transformation efficiency – cDNA or gDNA library construction – <i>Agrobacterium</i>-mediated transformation – Rifampicin resistance (pTiBo542DT-DNA)

GoldBio Competent Cells Characteristics and Applications



Products	Transformation Efficiency	Applications and Characteristics
<u>Auxo-Agro[®] EHA105 Chemically Competent Cells (CC-168)</u>	$\geq 1 \times 10^4$ CFU/ μ g	– High transformation efficiency – cDNA or gDNA library construction – <i>Agrobacterium</i>-mediated transformation – Rifampicin resistance (pTiBo542DT-DNA)
<u>EHA105 (pSoup) Agrobacterium ElectroCompetent Cells (CC-235)</u>	$\geq 4 \times 10^8$ CFU/ μ g	– High transformation efficiency – cDNA or gDNA library construction – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pEHA105(pTiBo542DT-DNA) – Rifampicin resistance (pTiBo542DT-DNA)
<u>EHA105 (pSoup) Agrobacterium Chemically Competent Cells (CC-118)</u>	$\geq 5 \times 10^3$ CFU/ μ g	– High transformation efficiency – cDNA or gDNA library construction – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pEHA105(pTiBo542DT-DNA) – Rifampicin resistance (pTiBo542DT-DNA)

GoldBio Competent Cells Characteristics and Applications



Products	Transformation Efficiency	Applications and Characteristics
<u>EHA105 (pSoup-P19) Agrobacterium ElectroCompetent Cells (CC-245)</u>	$\geq 3 \times 10^8$ CFU/ μ g	– High transformation efficiency – cDNA or gDNA library construction – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pEHA105(pTiBo542DT-DNA) – Rifampicin resistance (pTiBo542DT-DNA)
<u>EHA105 (pSoup-P19) Agrobacterium Chemically Competent Cells (CC-128)</u>	$\geq 2 \times 10^3$ CFU/ μ g	– High transformation efficiency – cDNA or gDNA library construction – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pEHA105(pTiBo542DT-DNA) – Rifampicin resistance (pTiBo542DT-DNA)
<u>EHA105 (pSuperAgro™ v4tet) Agrobacterium Electrocompetent Cells (CC-634)</u>	$\geq 1 \times 10^7$ CFU/ μ g	– High transformation efficiency – AcdS and GabT activity, driven by a single lac promotor – cDNA or gDNA library construction – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pEHA105(pTiBo542DT-DNA) – Rifampicin resistance (pTiBo542DT-DNA)

GoldBio Competent Cells Characteristics and Applications

Products	Transformation Efficiency	Applications and Characteristics
<u>EHA105 (pSuperAgro™ v4tet) Agrobacterium Chemically Competent Cells (CC-534)</u>	$\geq 1 \times 10^3$ CFU/ μ g	– High transformation efficiency – AcdS and GabT activity, driven by a single lac promotor – cDNA or gDNA library construction – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pEHA105(pTiBo542DT-DNA) – Rifampicin resistance (pTiBo542DT-DNA)
<u>GV3101 Agrobacterium Electrocompetent Cells (CC-207)</u>	$\geq 2 \times 10^8$ CFU/ μ g	– Very high transformation efficiency – cDNA or gDNA library construction – T-DNA binary system with the Ti plasmid pMP90(pTiC58DT-DNA) – Rifampicin resistance and gentamicin resistance – <i>Agrobacterium</i>-mediated transformation
<u>GV3101 Agrobacterium Chemically Competent Cells (CC-105)</u>	$\geq 6 \times 10^4$ CFU/ μ g	– Very high transformation efficiency – cDNA or gDNA library construction – T-DNA binary system with Ti plasmid pTiC58DT-DNA – Rifampicin resistance and gentamicin resistance – <i>Agrobacterium</i>-mediated transformation



GoldBio Competent Cells Characteristics and Applications

Products	Transformation Efficiency	Applications and Characteristics
<u>GV3101 (pSoup) Agrobacterium Electrocompetent Cells (CC-217)</u>	$\geq 8 \times 10^7$ CFU/ μ g	– Very high transformation efficiency – cDNA or gDNA library construction – T-DNA binary system with Ti plasmid pTiC58DT-DNA – Rifampicin resistance and gentamicin resistance – <i>Agrobacterium</i>-mediated transformation
<u>GV3101 (pSoup) Agrobacterium Chemically Competent Cells (CC-115)</u>	$\geq 1.5 \times 10^3$ CFU/ μ g	– Very high transformation efficiency – cDNA or gDNA library construction – T-DNA binary system with Ti plasmid pTiC58DT-DNA – Rifampicin resistance and gentamicin resistance – <i>Agrobacterium</i>-mediated transformation
<u>GV3101 (pSoup-P19) Agrobacterium Electrocompetent Cells (CC-227)</u>	$\geq 1 \times 10^8$ CFU/ μ g	– Very high transformation efficiency – cDNA or gDNA library construction – T-DNA binary system with Ti plasmid pMP90 (pTiC58DT-DNA) – Rifampicin resistance, tetracycline resistance and gentamicin resistance – <i>Agrobacterium</i>-mediated transformation

GoldBio Competent Cells Characteristics and Applications



Products	Transformation Efficiency	Applications and Characteristics
<u>GV3101 (pSoup-P19) Agrobacterium Chemically Competent Cells (CC-125)</u>	$\geq 3 \times 10^3$ CFU/ μ g	– Very high transformation efficiency – cDNA or gDNA library construction – T-DNA binary system with Ti plasmid pMP90 (pTiC58DT-DNA) – Rifampicin resistance, tetracycline resistance and gentamicin resistance – <i>Agrobacterium</i>-mediated transformation
<u>GV3101 (pSuperAgro™ v4tet) Agrobacterium Electrocompetent Cells (CC-614)</u>	$\geq 1 \times 10^7$ CFU/ μ g	– Very high transformation efficiency – AcdS and GabT activity, driven by a single lac promotor – cDNA or gDNA library construction – T-DNA binary system with Ti plasmid pMP90 (pTiC58DT-DNA) – Rifampicin resistance, tetracycline resistance and gentamicin resistance – <i>Agrobacterium</i>-mediated transformation
<u>GV3101 (pSuperAgro™ v4tet) Agrobacterium Chemically Competent Cells (CC-514)</u>	$\geq 1 \times 10^3$ CFU/ μ g	– Very high transformation efficiency – AcdS and GabT activity, driven by a single lac promotor – cDNA or gDNA library construction – T-DNA binary system with Ti plasmid pMP90 (pTiC58DT-DNA) – Rifampicin resistance, tetracycline resistance and gentamicin resistance – <i>Agrobacterium</i>-mediated transformation

GoldBio Competent Cells Characteristics and Applications



Products	Transformation Efficiency	Applications and Characteristics
<u>LBA4404 Agrobacterium Chemically Competent Cells (CC-107)</u>	$\geq 1 \times 10^4$ CFU/ μ g	– High transformation efficiency – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pAL4404 – Rifampicin resistance
<u>LBA4404 Agrobacterium Electrocompetent Cells (CC-220)</u>	$\geq 4 \times 10^6$ CFU/ μ g	– High transformation efficiency – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pAL4404 – Rifampicin resistance
<u>Auxo-Agro[®] LBA4404 Electrocompetent Cells (CC-267)</u>	$\geq 4 \times 10^6$ CFU/ μ g	– High transformation efficiency – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pAL4404 – Rifampicin resistance
<u>Auxo-Agro[®] LBA4404 Chemically Competent Cells (CC-167)</u>	$\geq 1 \times 10^4$ CFU/ μ g	– High transformation efficiency – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pAL4404 – Rifampicin resistance



GoldBio Competent Cells Characteristics and Applications

Products	Transformation Efficiency	Applications and Characteristics
<u>LBA4404 (pSuperAgro™ v4) Agrobacterium Electrocompetent Cells (CC-644)</u>	$\geq 1 \times 10^7$ CFU/ μ g	– High transformation efficiency – AcdS and GabT activity, driven by a single lac promotor – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pAL4404 – Rifampicin resistance
<u>LBA4404 (pSuperAgro™ v4) Agrobacterium Chemically Competent Cells (CC-544)</u>	$\geq 1 \times 10^3$ CFU/ μ g	– High transformation efficiency – AcdS and GabT activity, driven by a single lac promotor – <i>Agrobacterium</i>-mediated transformation – T-DNA binary system with the Ti plasmid pAL4404 – Rifampicin resistance