

Growth Factor Data Sheet

GoldBio growth factors are manufactured for **RESEARCH USE ONLY** and cannot be sold for human consumption!

CXCL12 is a CXC chemokine lacking the ELR motif. It was first found in bone marrow stromal cells and is expressed in a wide variety of cells with the exception blood cells. CXCL12 acts as a ligand for CXCR4. It is chemotactic for human hematopoietic progenitor cells expressing CD34, mononuclear cells, and cerebellar granule cells. CXCL12 plays a role in lymphopoiesis, myelopoiesis, vascularization of the gastrointestinal tract, and cerebellar development. CXCL12-beta is one of six alternatively spliced isoforms of the CXCL12 gene and is used as the canonical sequence.

Catalog Number	2110-12B
Product Name	CXCL12-beta, Human Recombinant Human Chemokine (C-X-C motif) Ligand 12, Beta (CXCL12 β) Stromal Cell-Derived Factor 1 (SDF1) Small Inducible Cytokine Subfamily B Member 12 (SCYB12) Pre-B Cell Growth-Stimulating Factor (PBSF) Thymic Lymphoma Cell-Stimulating Factor (TLSF)
Source	<i>Escherichia coli</i>
MW	~8.5 kDa (72 amino acids)
Sequence	KPVLSYRCP CRFFESHVAR ANVKHLKILN TPNCALQIVA RLKNNNRQVC IDPKLKWIQE YLEKALNKRF KM
Accession Number	P48061
Purity	>97% by SDS-PAGE and HPLC analyses
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using PHA and rHuIL-2 activated human peripheral blood T-lymphocytes is in a concentration range of 20-80 ng/ml.
Formulation	Sterile filtered white lyophilized powder. Purified and tested for use in cell culture.
Storage/Handling	This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage. The reconstituted sample can be apportioned into working aliquots and stored at -80 °C for up to 6 months. Avoid repeated freeze/thaw cycles.
Reconstitution	The sample should be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in a siliconized tube using PBS that contains a 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Reconstituted solutions are stable for up to one week at 2-8°C. Stock solutions should be aliquoted and stored at -80°C. Further dilutions should be made in appropriate buffered solutions containing BSA or serum.