

Growth Factor Data Sheet

GoldBio growth factors are manufactured for **RESEARCH USE ONLY** and cannot be sold for human consumption!

CXCL1 is an ELR type of CXC chemokine. It is secreted by human melanoma cells and expressed by neutrophils, monocytes, keratinocytes, macrophages, epithelial cells, and fibroblasts. CXCL1 participates in chemotaxis and cell activation of neutrophils. CXCR2 is the receptor that CXCL1 binds. CXCL1 inhibits embryonic oligodendrocyte precursor migration, suggesting a role in spinal cord development. It also plays a role in wound healing, angiogenesis, and tumorigenesis. CXCL1 is believed to be the major inflammatory factor in recurrent herpetic stromal keratitis.

Catalog Number	2110-01
Product Name	CXCL1, Human Recombinant Human Chemokine (C-X-C motif) Ligand 1 Melanoma Growth Stimulating Activity, Alpha (MGSA, MGSA-a) Fibroblast Secretory Protein (FSP) Growth Related Oncogene 1 (GRO1, GROa) Neutrophil-Activating Protein 3 (NAP3, NAP-3) Small Inducible Cytokine Subfamily B Member 1 (SCYB1)
Source	<i>Escherichia coli</i>
MW	~7.9 kDa (73 amino acids)
Sequence	ASVATELRQC CLQTLQGIHP KNIQSVNVKS PGPHCQAQTEV IATLKNGRKA CLNPASPIVK KIIKMLNSD KSN
Accession Number	P09341
Purity	>97% by SDS-PAGE and HPLC analyses
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human peripheral blood neutrophils is in a concentration range of 10-50 ng/ml.
Formulation	Sterile filtered white lyophilized powder. Purified and tested for use in cell culture.
Storage/Handling	This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage. The reconstituted sample can be apportioned into working aliquots and stored at -80 °C for up to 6 months. Avoid repeated freeze/thaw cycles.
Reconstitution	The sample should be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in a siliconized tube using PBS that contains a 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Reconstituted solutions are stable for up to one week at 2-8°C. Stock solutions should be aliquoted and stored at -80°C. Further dilutions should be made in appropriate buffered solutions containing BSA or serum.