

Growth Factor Data Sheet

GoldBio growth factors are manufactured for **RESEARCH USE ONLY** and cannot be sold for human consumption!

IL2 is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. It is expressed by CD4+ and CD8+ T-cells, $\gamma\delta$ T-cells, B-cells, dendritic cells, and eosinophils. The receptor for IL2 consists of three subunits that are present on the cell surface in varying preformed complexes. The 55 kDa IL2 R α is specific for IL2 and binds with low affinity. The 75 kDa IL2 R β , which is also a component of the IL15 receptor, binds IL2 with intermediate affinity. The 64 kDa common gamma chain γ_c /IL2 R γ , which is shared with the receptors for IL4, IL7, IL9, IL15 and IL21, does not interact independently with IL2. Upon ligand binding, signal transduction is performed by both IL2 R β and γ_c . It drives resting T-cells to proliferate and induces IL2 and IL2 R α synthesis. It contributes to T-cell homeostasis by promoting the Fas-induced death of naive CD4+ T-cells but not activated CD4+ memory lymphocytes. IL2 plays a central role in the expansion and maintenance of regulatory T-cells, although it inhibits the development of Th17 polarized cells. This recombinant IL2 has been modified by substituting a serine for cysteine at position 141 of the original sequence.

Catalog Number	1010-02A
Product Name	IL2 (Cys141Ser), Equine Recombinant Equine Interleukin 2 (Cys141Ser) IL-2, IL2 T-cell Growth Factor (TCGF)
Source	<i>Escherichia coli</i>
MW	~14.9 kDa (129 amino acid chains)
Sequence	APTSSSKRET QQQLKQLQMD LKLLLEGVNN NKNPKLSKML TFKINMPKKA TELKHLQCLE EELKPLEEML KNFLSKDIKE LMSNINVTVL GLKGSETRFT CEYDDEGTI VEFLNKWITF SQSIFSTMT
Accession Number	P37997
Purity	>97% by SDS-PAGE and HPLC analyses
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using murine CTLL-2 cells is less than 1.0 μ g/ml, corresponding to a specific activity of >1000 IU/mg.
Formulation	Sterile filtered white lyophilized powder. Purified and tested for use in cell culture.
Storage/Handling	This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage. The reconstituted sample can be apportioned into working aliquots and stored at -80 °C for up to 6 months. Avoid repeated freeze/thaw cycles.
Reconstitution	The sample should be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in a siliconized tube using PBS that contains a 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Reconstituted solutions are stable for up to one week at 2-8°C. Stock solutions should be aliquoted and stored at -80°C. Further dilutions should be made in appropriate buffered solutions containing BSA or serum.